

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/338147752>

# Using Connected Learning Design Principles to Further Co-Create a Critical Speech Therapy Game

Conference Paper · December 2019

CITATIONS

0

READS

112

5 authors, including:



Jared Duval

University of California, Santa Cruz

12 PUBLICATIONS 26 CITATIONS

[SEE PROFILE](#)



Elena Márquez Segura

University of California, Santa Cruz

65 PUBLICATIONS 592 CITATIONS

[SEE PROFILE](#)



Elizabeth Jessica Goldman

University of California, Santa Cruz

5 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



All the Feels [View project](#)



The ExerCube – Towards Designing and Evaluating Attractive and Effective Fitness Game Environments [View project](#)

# Using Connected Learning Design Principles to Further Co-Create a Critical Speech Therapy Game

Jared Duval, University of California Santa Cruz, [JDuval@ucsc.edu](mailto:JDuval@ucsc.edu)

Elena Márquez Segura, University of California Santa Cruz, [elena.marqz@gmail.com](mailto:elena.marqz@gmail.com)

Elizabeth Goldman, University of California Santa Cruz, [elijgoldm@ucsc.edu](mailto:elijgoldm@ucsc.edu)

Su-hua Wang, University of California Santa Cruz, [suhua@ucsc.edu](mailto:suhua@ucsc.edu)

Sri Kurniawan, University of California Santa Cruz, [Skurnia@ucsc.edu](mailto:Skurnia@ucsc.edu)

**Abstract:** Therapy can be costly, time-consuming, repetitive, and difficult. Games have the power to teach transferable skills, can turn repetitive tasks into engaging mechanics, have been proven to be effective at delivering various forms of therapy, and can be deployed at large scales. Therapy games represent fertile ground for connected learning. In this work, we collaborate with 7 children with corrected cleft palate aged 2-10 and their parents during their yearly visit to UC Davis Medical center to co-create and evaluate Spokelt, a speech therapy game. Each of these children come from low socioeconomic statuses with limited access to speech therapy and would benefit from the amplified opportunities of new media in connected learning. Throughout the study, we ran multiple cascading participatory design sessions using design principles of connected learning, which culminated in the design of two new medium-fidelity prototypes presented in this paper.

## Introduction

In 2012, nearly 8% of children aged 3-17 in the United States had a communication disorder, and younger children, boys, and non-Hispanic white children were more likely than other children to receive an intervention service for their disorder (*National Center for Health Statistics, 2015*). Correcting speech is an important issue because children with speech impairments have high risks of behavioral problems and increased symptoms of depression (Hunt, Burden, Hepper, Stevenson, & Johnston, 2007). They show more deficits in social and academic competencies, score higher for social problems (Feragen, Kvaalem, Rumsey, & Borge, 2010), and are more likely to be teased in social settings (Hunt et al., 2007). These are serious consequences for a skill that can be improved (Aten et al., 1982; Johnson & Pring, 1990; Robertson & Thomson, 1984). Even those who undergo corrective surgery tend to display a delay in scholarship, have a lower income, marry later in life and become independent from their parents significantly later (Sousa, Devare, & Ghanshani, 2009). After surgery, practice and support are the keys to improving speech (Cicerone et al., 2005). While Speech-Language Pathologists (SLPs) employ many engaging games (Mashima & Doarn, 2008), progress in speech is often slow or lost at home because speech practice at home is usually hindered by a lack of intrinsic motivation due to the tedious and repetitive idiosyncratic nature of traditional speech therapy curriculums (Aten, Caligiuri, & Holland, 1982). Parents also experience significant difficulties prompting their children to complete speech exercises at home, especially if they are very young (Miesenberger, Klaus, Zagler, & Karshmer, 2010).

Speech therapy games could help children practice at home and expedite their recovery (Rubin, 2017). Games have the potential to turn repetition, such as repeating words in speech therapy, into an element that is recognized as useful for progress in the game (Kaufman, 2010). Literature had shown that games are an effective educational intervention and medium to convey and support feelings of self-efficacy due to the immersive and pervasive virtual environment and that it had been shown to work effectively as educational interventions (Gee, 2005). Games have demonstrated the ability to teach while providing a motivating and interactive environment (Virvou, Katsionis, & Manos, 2005), are effective with children (Rosas et al., 2003) and can be as effective as face-to-face instruction (Randel, Morris, Wetzel, & Whitehill, 1992). A variety of serious games for health have been documented to be useful for their target populations, and are wide-ranging in their platforms, health outcomes and target populations, from an exergame to help blind children with balance (Morelli, Lieberman, Foley, & Folmer, 2014) to embodied persuasive games for adults in wheelchair (Gerling, Hicks, Kalyn, Evans, & Linehan, 2016) to mobile games for motivating tobacco-free life in early adolescence (Parisod et al., 2017).

## Background

Spokelt is a speech therapy game equipped with a critical speech recognition system capable of “hearing” many common speech errors. Spokelt contains numerous mini-games that target specific types of speech therapy, such as rhythm and articulation, that fit within an overarching narrative. Spokelt’s narrative, mechanics, and characters were iteratively co-designed and presented in our previous works (J. Duval, 2017; J. S. Duval, Márquez Segura, & Kurniawan, 2018).

## Method

We visited our collaborators at UC Davis Medical Center where children with corrected cleft are assessed by different medical experts including speech pathologists, behavioral therapists, plastic surgeons, sleep therapists, dentists, etc. We were invited to take part in a “clinical day” where patients are assigned to a room while a variety of doctors rotate in and out to see each patient. Per request of the speech language pathologist and in order to minimize interference to the doctors’ rotations, we designed co-creation sessions that lasted 10 minutes, which would be conducted during “gaps” where no medical professionals were present in the patient’s assigned room. Seven children participated in our study whose ages ranged from 2 years, 7 months to 10 years, 9 months ( $M = 6$  years 1 month,  $SD = 3$  years 6 months). They were accompanied by their parent(s), who observed them play and were also interviewed. We also interviewed 2 speech language pathologists. In all, we collected data on 7 children, 7 parents, and 2 Speech Language Pathologists.

## Connected Learning Design Principals for Meaningful Co-Creation

We visited our collaborators at UC Davis Medical Center, where children with a corrected cleft are assessed by different medical experts, including speech pathologists, behavioral therapists, plastic surgeons, sleep therapists, dentists, etc. We were invited to take part in a “clinical day” where patients are assigned to a room while a variety of doctors rotate in and out to see each patient. Per the request of the speech-language pathologist and in order to minimize interference to the doctors’ rotations, we designed co-creation sessions that lasted 10 minutes, which would be conducted during “gaps” where no medical professionals were present in the patient’s assigned room. Seven children participated in our study whose ages ranged from 2 years, 7 months to 10 years, 9 months ( $M = 6$  years 1 month,  $SD = 3$  years 6 months). They were accompanied by their parent(s), who observed them play and were also interviewed. We also interviewed 2 speech-language pathologists. In all, we collected data on 7 children, 7 parents, and 2 Speech-Language Pathologists.



Figure 1. Design probes used for cascading participatory design sessions

## Everything is interconnected

In order to come up with play scenarios, children were provided with tangible co-design probes, toys, and props (see Figure 1). The tangible representations of our characters connected the play session to the Spokelt Universe. The animal flashcards and word flashcards were chosen because they are rated as developmentally appropriate words for ages 3 and higher. The letter and number magnets were useful random character or number generators. The stickers were a gift to the children for participating. At the beginning of the session, we played Spokelt's cinematic on a laptop to remind participants of the story and to help them connect the virtual and tangible characters. Next, children were presented with the rest of the tangible co-design probes to create and play game scenarios. In between sessions, we created medium-fidelity prototypes of these play scenarios in Adobe XD. We then had the following participants play and critique the adobe XD prototypes so that each design was tested and iterated with subsequent participants, hence cascading participatory design sessions.

## Learning Happens by Doing

To inspire them, they were asked first to play either a play scenario we had prepared or one that a previous child had proposed. Children were encouraged to iterate these scenarios, changing the rules, objects used, etc. and to play the new scenario. For example, if a game required them to repeat a word X number of times, where X is a number magnet pulled from a container, we asked that they physically draw a number and repeat the word that many times.

## Challenge is Constant

In the latter cascading co-creation sessions, the children played scenarios that had already been mocked up in Adobe XD. Facilitators presented the iPad with the prototype running and read the written instructions out loud to the children. Using Wizard of Oz techniques (Dow et al., 2005; Fraser & Gilbert, 1991), facilitators took the role of the speech recognition system and the narrator. They would clarify instructions, give helpful hints, and provide feedback when necessary as the game would do when it is finished. We took note of when they were engaged, which design elements, the child seemed to enjoy the most, and when the child needed more support.

## Anyone Can Participate

After playing the previously made designs, we invited children to come up with another play scenario using the tangible probes. The facilitator helped the children polish it with questions such as: what happens next? How do they do that? What do they need to say to help? The facilitator reminded players that all of the challenges should require speech. Once the play scenario was concluded, the child played it once or twice with the help of the facilitators. Last, the children were asked general questions about the play scenario, such as: what do you like the most about this game? Why? What do you like the least? Would you like playing this game at home? They were also asked which game of those they had played they liked the most. At the end of the session, the child chose a game character sticker to take home as a souvenir.

## Designs




Figure 2. Adobe XD medium fidelity prototype of farm-to-table mini-game rapidly developed during gaps

The first design (see Figure 2) is a farm-to-table mini-game where players cultivate the crops needed to make a salad. In this mini-game, players pick the various fruits and vegetables by saying the name of the vegetable. If the salad requires five tomatoes, they would say tomato five times. During the Wizard of Oz Adobe XD playtests, we gained valuable knowledge about how we naturally support children and ideas on how this should be incorporated into the final product: It helps to repeat the same directions but worded in different ways. Different children can complete the same challenge in various directions. Silence and numerous incorrect attempts indicate confusion. We found the magic number to be three seconds of silence or 3 incorrect attempts in a row indicates the child might need support. If there is a list of items required, it helps to provide example solutions. The game should ask the child if they need help when they might be struggling. Sometimes they need more time to complete the task or a few more tries. We also found that children enjoy silly euphonic words like “pickle,” which may be more challenging to guess, but are more rewarding because they are fun to say.



Figure 3. Adobe XD medium fidelity prototype of farm animals mini-game rapidly developed during gaps

The second design (see Figure 3) started as a predesigned scenario to spur creativity. However, the children enjoyed playing it and iterating on it, so we decided to mock it up in Adobe XD on the scene. The characters were at a farm with no animals. To bring an animal to the farm, the child needed to say the animal's name and the sound that the animal makes. To give the animal a name, they needed to say a name that started with the same sound as the type of animal. At this point, the facilitator asked the child to pick up a number card out of a box randomly. “[X] times! You need to say their name [X] times, so they remember it”. When the child calls the [animal] [X] times, that animal’s card is turned over. The same process is repeated with the 3 other animals.

## Discussion

In this paper, we present a novel co-design method that emerged out of the constraints and possibilities of our testing environment: a tight fast-past cascading co-design protocol featuring play, tangibles, on-scene medium-fidelity prototyping, Wizard of Oz techniques, and continued iteration of designs using the design principles of connected learning:

### Learning Happens by Doing

We found two aspects essential: First, a warm-up scenario is helpful so that children understand what they need to create. Second, having tangible play props. The character props allowed children to immerse themselves in the universe and come up with interesting play scenarios. Children liked these

tangible characters (even wanted to keep them), held onto them, played with them, and felt comfortable using them to make new games.

### **Challenge is Constant**

The props directly influenced emerging games, which is why choosing appropriate props was extremely important. We used wooden magnets to act as random letters and number generators, age-appropriate flashcards to prompt words that need to be spoken, and tangible felted characters crafted to look like the characters in the game. The props directly influence resulting designs, so if the facilitators are not careful with these choices, the emerging play may be constrained by rules that do not fit goals. It can be challenging to redirect focus and play to the goals of the session. Our probes, especially the flashcards, helped keep play constrained to using speech. Because children had already played the game, they understood the constraints of the Spokelt Universe. It helps to define the play space clearly.

### **Anyone Can Participate**

Co-creating a game—even if it is a sketch of a play scenario— is challenging, and even more so if the co-creators do not have a background in game design and are children (!). However, children are experts in make-believe, and crafting a magic circle comes naturally to them. The challenge is crafting “the right circle,” one that can help them in their therapy.

### **Everything is Interconnected**

We used breaks between sessions to create medium-fidelity interactive prototypes on-scene based on games children made with our tangible probes. This meant valuable design knowledge was not lost to time because it was made immediately after sessions. Each design could be iterated multiple times in the day, with multiple children, in a medium-fidelity environment (Adobe XD). Children evaluated other children’s designs, which gave insights into which features were essential to multiple children versus just the child who made the game.

### **References**

- Aten, J. L., Caligiuri, M. P., & Holland, A. L. (1982). The efficacy of functional communication therapy for chronic aphasic patients. *Journal of Speech and Hearing Disorders*, 47(1), 93–96.
- Cicerone, K. D., Dahlberg, C., Malec, J. F., Langenbahn, D. M., Felicetti, T., Kneipp, S., ... Others. (2005). Evidence-based cognitive rehabilitation: updated review of the literature from 1998 through 2002. *Archives of Physical Medicine and Rehabilitation*, 86(8), 1681–1692.
- Dow, S., MacIntyre, B., Lee, J., Oezbek, C., Bolter, J. D., & Gandy, M. (2005). Wizard of Oz support throughout an iterative design process. *IEEE Pervasive Computing*, 4(4), 18–26.
- Duval, J. (2017). A mobile game system for improving the speech therapy experience. In *Proceedings of the 19th International Conference on Human-Computer Interaction with Mobile Devices and Services* (p. 72). ACM.
- Duval, J. S., Márquez Segura, E., & Kurniawan, S. (2018). Spokelt: A Co-Created Speech Therapy Experience. In *Extended Abstracts of the 2018 CHI Conference on Human Factors in Computing Systems* (p. D501). ACM.

- Feragen, K. B., Kvalem, I. L., Rumsey, N., & Borge, A. I. (2010). Adolescents with and without a facial difference: the role of friendships and social acceptance in perceptions of appearance and emotional resilience. *Body Image, 7*(4), 271–279.
- Fraser, N. M., & Gilbert, G. N. (1991). Simulating speech systems. *Computer Speech & Language, 5*(1), 81–99.
- Gee, J. P. (2005). Learning by design: Good video games as learning machines. *E-Learning and Digital Media, 2*(1), 5–16.
- Gerling, K., Hicks, K., Kalyn, M., Evans, A., & Linehan, C. (2016). Designing movement-based play with young people using powered wheelchairs. In *Proceedings of the 2016 CHI Conference on Human Factors in Computing Systems* (pp. 4447–4458). ACM.
- Hunt, D. O., Burden, D. D., Hepper, D. P., Stevenson, D. M., & Johnston, D. C. (2007). Parent Reports of the Psychosocial Functioning of Children with Cleft Lip and/or Palate. *The Cleft Palate-Craniofacial Journal, 44*(3), 304–311. <https://doi.org/10.1597/05-205>
- Ito, M., Gutiérrez, K., Livingstone, S., Penuel, B., Rhodes, J., Salen, K., ... Watkins, S. C. (2013). *Connected learning: An agenda for research and design*. BookBaby.
- Johnson, J. A., & Pring, T. R. (1990). Speech therapy and Parkinson's disease: a review and further data. *British Journal of Disorders of Communication, 25*(2), 183–194.
- Kaufman, D. (2010). *Educational Gameplay and Simulation Environments: Case Studies and Lessons Learned: Case Studies and Lessons Learned*. IGI Global.
- Márquez Segura, E., Waern, A., Márquez Segura, L., & López Recio, D. (2016). Playification: The PhySeEar case. In *Proceedings of the 2016 Annual Symposium on Computer-Human Interaction in Play* (pp. 376–388). ACM.
- Mashima, P. A., & Doarn, C. R. (2008). Overview of telehealth activities in speech-language pathology. *Telemedicine and E-Health, 14*(10), 1101–1117.
- McGonigal, J. (2011a). *Reality is broken: Why games make us better and how they can change the world*. Penguin.

- McGonigal, J. (2011b). *We Don't Need No Stinkin' Badges: How to Re-invent Reality Without Gamification*. Presented at the Games Developers Conference, GDC. Retrieved from <http://www.gdcvault.com/play/1014576/We-Don-t-Need-No>
- Miesenberger, K., Klaus, J., Zagler, W., & Karshmer, A. (2010). *Computers Helping People with Special Needs, Part II: 12th International Conference, ICCHP 2010, Vienna, Austria, July 14-16, 2010. Proceedings*. Springer Berlin Heidelberg. Retrieved from <https://books.google.com/books?id=jlbX8RxAf7UC>
- Morelli, T., Lieberman, L., Foley, J., & Folmer, E. (2014). An exergame to improve balance in children who are blind. In *FDG*. National Center for Health Statistics. (2015). Centers for Disease Control and Prevention. Retrieved from <https://www.cdc.gov/nchs/products/databriefs/db205.htm>
- Nicholson, S. (2015). A recipe for meaningful gamification. In *Gamification in education and business* (pp. 1–20).
- Parisod, H., Pakarinen, A., Axelin, A., Danielsson-Ojala, R., Smed, J., & Salanterä, S. (2017). Designing a Health-Game Intervention Supporting Health Literacy and a Tobacco-Free Life in Early Adolescence. *Games for Health Journal*.
- Randel, J. M., Morris, B. A., Wetzel, C. D., & Whitehill, B. V. (1992). The effectiveness of games for educational purposes: A review of recent research. *Simulation & Gaming*, 23(3), 261–276.
- Robertson, S. J., & Thomson, F. (1984). Speech therapy in Parkinson's disease: a study of the efficacy and long term effects of intensive treatment. *International Journal of Language & Communication Disorders*, 19(3), 213–224.
- Rosas, R., Nussbaum, M., Cumsille, P., Marianov, V., Correa, M., Flores, P., ... Others. (2003). Beyond Nintendo: design and assessment of educational video games for first and second grade students. *Computers & Education*, 40(1), 71–94.
- Rubin, Z. (2017). *Development and evaluation of software tools for speech therapy* (PhD Thesis). University of California, Santa Cruz.



Sousa, A. D., Devare, S., & Ghanshani, J. (2009). Psychological issues in cleft lip and cleft palate.

*Journal of Indian Association of Pediatric Surgeons*, 14(2), 55–58. <https://doi.org/10.4103/0971-9261.55152>

Virvou, M., Katsionis, G., & Manos, K. (2005). Combining software games with education: Evaluation of its educational effectiveness. *Educational Technology & Society*, 8(2), 54–65.